

Galaxy Disks and Disk Galaxies
ASP Conference Series, Vol. 3 × 10⁸, 2000
J.G. Funes, S.J., and E.M. Corsini, eds.

SuperMassive Black Holes from the SUNNS survey

Marc Sarzi

*Dipartimento di Astronomia, Università di Padova, Vicolo
dell'Osservatorio 5, I-35122 Padova, Italy*

Hans-Walter Rix

*Max-Planck-Institut für Astronomie, Königstuhl 17, Heidelberg,
D-69117, Germany*

Joseph C. Shields

Physics & Astronomy Department, Ohio University, Athens, OH 45701

Gregory Rudnick, Daniel H. McIntosh

Steward Observatory, University of Arizona, Tucson, AZ 85721

Luis C. Ho

*The Observatories of the Carnegie Institution of Washington, 813 Santa
Barbara St., Pasadena, CA 91101-1292*

Alexei V. Filippenko

*Astronomy Department, University of California, Berkeley, CA
94720-3411*

Wallace L. W. Sargent

Palomar Observatory, Caltech 105-24, Pasadena, CA 91125

Over the last few years evidence has mounted that supermassive black holes (SMBH) are nearly ubiquitous in galactic centers (see, e.g., Ho 1999 for a review). The possibility that a close connection between the formation and evolution of SMBHs and of their host galaxies exists, has recently been bolstered by reports of a correlation between the black-hole mass $M_{\bullet}$ and the velocity dispersion of the stellar bulge σ (Ferrarese & Merritt 2000; Gebhardt et al. 2000; Merritt & Ferrarese 2000). Despite this recent progress, the demography of SMBHs is far from complete. Most of the SMBHs have been detected in bulge-dominated systems, and it is not clear if the $M_{\bullet}$ - σ relation holds also in the case of disk-dominated galaxies, or when a bar is present.

In order to address this and other issues we initiated a Survey of Nearby Nuclei with STIS (SUNNS) on *HST*, with kinematic information obtained from measurements of nebular emission lines for 24 galaxies. From this starting sample, four galaxies displayed symmetric gas velocity curves that could be consistently modeled as a rotating disk in the joint potential of the stellar bulge and a putative central black hole. An extensive description of the analysis for these galaxies




Figure 1. *Left:* Our 3 $M_{\bullet}$ determinations (open squares) along with some of the $M_{\bullet}$ measurements from which the Merritt & Ferrarese (2000) relation (solid line) was derived ($M_{\bullet}$ from reverberation mapping are not shown). Different symbols correspond to the different tracer of the potential used to obtain $M_{\bullet}$. *Right:* Same as *left* except that different symbols now indicate the host galaxy Hubble-type.

can be found in Sarzi et al. (2000). In Figure 1 we show how the three sources for which we obtained well-determined values of $M_{\bullet}$, namely NGC 2787(*SB0*), NGC 4459(*S0*), NGC 4596(*SB0*), fall in the $M_{\bullet}$ - σ plane.

Reliable $M_{\bullet}$ determinations based on gas kinematics have been limited so far to ionized gas measurements in ellipticals and water-maser measurements in late-type spirals. Our $M_{\bullet}$ measurements fill the corresponding gap in the $M_{\bullet}$ - σ plane, and our SUNNS survey illustrates both the difficulty in finding regular gas motions in disk galaxies and the need for an accurate preselection of the targets. Finally our $M_{\bullet}$ determinations are not only consistent with the $M_{\bullet}$ - σ relation, but also reinforce it. Indeed the number of disk galaxies with accurate $M_{\bullet}$ measurements (excluding reverberation mapping measurements) has now been increased from 9 to 12, the number of S0's from 5 to 8, and the number of barred galaxies has been doubled.

References

- Ferrarese, L. & Merritt, D. 2000, ApJ, 539, L9
 Gebhardt, K. et al. 2000, ApJ, 539, L13
 Merritt, D. & Ferrarese, L., ApJ, in press, astro-ph/0008310
 Ho, L. 1999, in Observational Evidence for the Black Holes in the Universe, ed S.K. Chakrabarti (Dordrecht: Kluwer), 157
 Sarzi, M. et al. 2000, ApJ, in press, astro-ph/0010240